L11 ANSWER 93 OF 95 DGENE COPYRIGHT 2006 The Thomson Corp on STN

ACCESSION NUMBER: AAW34569 peptide DGENE

TITLE: Recovery of fibrinogen using polysaccharide solid support

coupled to fibrinogen-binding peptide - requires only mild

elution buffers

INVENTOR: Goss N; Kanellos J; Oates A; Pham H

PATENT ASSIGNEE: (CSLC-N)CSL LTD.

PATENT INFO: WO 9726280 A1 19970724 24

APPLICATION INFO: WO 1997-AU13 19970114 PRIORITY INFO: AU 1996-7564 19960116

DOCUMENT TYPE: Patent LANGUAGE: English

AB

OTHER SOURCE: 1997-385298 [35]

DESCRIPTION: Synthetic fibrinogen binding peptide 2.

AB Peptides AAW34568-71 are synthetic fibrinogen binding peptides.

A tripeptide, Gly-Pro-Arg is also claimed.

This tripeptide sequence corresponds to the first 3 amino acids of the alpha-chain exposed by the thrombin catalysed release of the fibrinopeptide A in all vetebrate species. In peptide AAW34568, the addition of a proline residue at position 4 increases the affinity of the peptide for fibrinogen almost tenfold. These synthetic fibrinogen binding peptides are immobilised on a novel polysaccharide support (e.g. Sephadex), to which they are coupled through a spacer or linker moiety. This linker moiety comprises a chain of greater then 7 atoms. The solid support is useful for the recovery and isolation of fibrinogen from material such as plasma, plasma fractions and fibrinogen-containing cell culture media arising from the production of fibrinogen by recombinant DNA techniques. The process is superior to other known affinity isolation procedures in that only mild elution buffers are required to recover the bound fibrinogen.

Peptides AAW34568-71 are synthetic fibrinogen binding peptides.

A tripeptide, Gly-Pro-Arg is also claimed. This tripeptide sequence corresponds to the first 3 amino acids of the alpha-chain exposed by the thrombin catalysed release of the fibrinopeptide A in all vetebrate species. In peptide AAW34568, the addition of a proline residue at position 4 increases the affinity of the peptide for fibrinogen almost tenfold. These synthetic fibrinogen binding peptides are immobilised on a novel polysaccharide support (e.g. Sephadex), to which they are coupled through a spacer or linker moiety. This linker moiety comprises a chain of greater then 7 atoms. The solid support is useful for the recovery and isolation of fibrinogen from material such as plasma, plasma fractions and fibrinogen-containing cell culture media arising from the production of fibrinogen by recombinant DNA techniques. The process is superior to other known affinity isolation procedures in that only mild elution buffers are required to recover the bound fibrinogen.

(FILE 'HOME' ENTERED AT 08:40:11 ON 20 FEB 2006)

FILE 'DISSABS, 1MOBILITY, AGRICOLA, AQUASCI, BIOTECHNO, COMPENDEX, COMPUAB, CONF, CONFSCI, ELCOM, HEALSAFE, IMSDRUGCONF, LIFESCI, OCEAN, PAPERCHEM2, PASCAL, POLLUAB, SOLIDSTATE, ADISCTI, ADISINSIGHT, ADISNEWS, ANABSTR, ANTE, AQUALINE, BIOENG, BIOSIS, ...' ENTERED AT 08:40:22 ON 20 FEB 2006

	1 2 2 2 0 0 0	
		E RICHTER?/AU
L:	5396	S E2
		E KOLL ROBERT?/AU
L	2 6	S E1 OR E2
		E BIEBER FRANZ?/AU
L3	3 28	S E1 OR E2
		E TSCHOPE?/AU
L	191	S E2
L5	22046	S GLY (A) PRO (A) ARG (A) PRO
Le	5 3	S L5 AND (L1 OR L2 OR L3 OR L4)
L'	7 3218	S GLY (A) PRO (A) ARG (A) PRO (A) LYS
L8	3 22046	S GLY (A) PRO (A) ARG
L9	781	S (L5 OR L7 OR L8) (S) (ADSORB? OR AFFINITY OR COUPLED OR
L:	10 162	S L9 (S) (FIBRIN?)
L	11 95	DUP REM L10 (67 DUPLICATES REMOVED)